

Fermi-surface properties of the perovskite superconductors

Claudius Gros, Roser Valentí

*Institut für Physik, Universität Dortmund,
44221 Dortmund, Germany,
email: UPH301 and UPH084 at DDOHRZ11*

We calculate Fermi-surface properties of the Cuprate superconductors within the three-band Hubbard model using a cluster expansion for the proper self-energy. The Fermi-surface topology is in agreement with angular-resolved photoemission data for dopings $\sim 20\%$. We discuss possible violations of the Luttinger sum-rule for smaller dopings and the role of van-Hove singularities in the density of states of the Zhang-Rice singlets. We calculate the shift in the chemical potential upon doping and find quantitative agreement with recent experiments.

PACS. 71.10 Many-body theory – electron states

PACS. 74.20 Hight- T_C – theory

PACS. 79.60 Photoemission

Introduction In the last years, the experimental characterization of Fermi-surface properties of the high-temperature superconductors has progressed with the help of angle-resolved^(1,2,3,4,5) and angle-integrated⁽⁶⁾ photoemission studies. The such determined Fermi-surface topology agrees quite well with results from band-structure calculations based on the local-density approximation⁽⁷⁾. This measured Fermi-surface topology has been taken as an input for theoretical calculations⁽⁸⁾ in the so called “Fermiology” scenario. Many theorists use, on the other hand, a simplified three-band Hamiltonian (also called Emery model⁽⁹⁾), which incorporates strong, local correlations, as exemplified by the composite-operator approach⁽¹⁰⁾. Determination of the Fermi-surface properties of the Emery model is of central interest and has been investigated in a series of recent studies (11,12,13,14).

It is difficult to calculate with numerical methods, like exact diagonalization^(15,16) or Quantum Monte-Carlo⁽¹⁷⁾, the Fermi-surface topology for the three-band model directly, due to cluster-size limitations. The numerical methods have been shown, on the other side, to be very useful for extracting the energy scales of the local charge- and spin excitations⁽¹⁵⁾. Here we propose that the *cluster self-energy* provides the missing link between such numerical approaches and calculation of momentum-dependent properties, like the Fermi-surface topology.

Method The self-energy of finite clusters with open boundary conditions contains in a very precise way information upon the Green’s function and self-energy of the *extended* system⁽¹⁸⁾. Let us consider the real-space components of the self-energy for both the extended system, $\Sigma_{i,j}(\omega)$, and for a given finite cluster with open boundary conditions, $\Sigma_{i,j}^{(c)}(\omega)$. It is straightforward, that all irreducible diagrams contributing to $\Sigma_{i,j}^{(c)}(\omega)$ are a subset of the diagrams contributing to $\Sigma_{i,j}(\omega)$. Therefore the cluster self-energy is a *systematic* approximation to the self-energy of the full system.

Typically one first evaluates the cluster Green’s function, $G_{i,j}^{(c)}(\omega)$, in an exact diagonalization study. The cluster self-energy is then obtained by inverting the cluster Dyson’s equation

$$G_{i,j}^{(c)}(\omega) = G_{i,j}^{(c,0)}(\omega) + \sum_{m,n} G_{i,n}^{(c,0)}(\omega) \Sigma_{m,n}^{(c)}(\omega) G_{n,j}^{(c)}(\omega), \quad (1)$$

where the i, m, n, j are sites of the cluster and $G_{i,j}^{(c,0)}(\omega)$ is the non-interacting cluster Green’s function.

The Green’s function of the full CuO₂ lattice is determined by inserting the $\Sigma_{m,n}^{(c)}(\omega)$ determined from Eq. (1), into the Dyson’s equation of the extended system

$$G_{i,j}(\omega) = G_{i,j}^{(0)}(\omega) + \sum_{m,n} G_{i,n}^{(0)}(\omega) \Sigma_{m,n}^{(c)}(\omega) G_{n,j}(\omega), \quad (2)$$

where the i, m, n, j run over all sites of the CuO₂ lattice and $G_{i,j}^{(0)}(\omega)$ is the non-interacting Green’s function of the CuO₂ plane. The Green’s function, $G_{i,j}(\omega)$, obtained

by Eq. (2) can be systematically improved considering larger and larger clusters in Eq. (1). The smallest cluster one could possibly consider is the one-site-cluster and Eq. (2) together with Eq. (1) would then reduce to the Hubbard-I approximation⁽¹⁹⁾. For the Emery model we consider the CuO_4 cluster for Eq. (1). Within this approximation both the important local charge fluctuations and the singlet-triplet splitting are included.

The passage of the quasi-particle peak through the chemical potential, determined from the $\mathbf{k}$ -dependent Green's function, $G_{\mathbf{k}}(\omega) \equiv 1/L \sum_{i,j} \exp[i\mathbf{k}(\mathbf{R}_i - \mathbf{R}_j)] G_{i,j}(\omega)$ determines then the Fermi-surface. The density of electrons is given by

$$n \equiv 2 \int_{-\infty}^{\infty} d\omega \left(\frac{-1}{\pi} \right) \text{Im} G(\omega) n_F(\omega - \mu) = 2 \int_{\mathbf{k} \in FS} \frac{d^2 k}{(2\pi)^2} Z_{\mathbf{k}}, \quad (3)$$

with $n_F(\omega - \mu)$ and $Z_{\mathbf{k}}$ being the Fermi distribution function and the quasiparticle spectral weight respectively. The factor of two in Eq. (3) comes from spin-degeneracy and the $\mathbf{k}$ -sum is over all occupied states. The density of electrons, as given by Eq. (3), is to be determined self-consistently⁽¹⁸⁾ as a function of the chemical potential, μ , by demanding it to be equal to the average density of electrons of the CuO_4 cluster, for which $\Sigma_{i,j}^{(c)}(\omega)$ is determined via Eq. (1).

Results For the three-band Hamiltonian we have chosen (in electron notation), following Eskes and Sawatzky⁽¹⁵⁾, $\epsilon_p - \epsilon_d = 5.3\text{eV}$ for the difference in oxygen- and copper orbital energies⁽²⁰⁾, $t_{dp} = 1.3\text{eV}$ and $t_{pp} = 0.65\text{eV}$ for the Cu-O and the O-O hopping matrix element respectively and $U_d = 8.8\text{eV}$ for the onsite Coulomb repulsion on the copper site.

From the Green's function of a CuO_4 cluster we have extracted the self-energy and used it as an approximation for the self-energy of the infinite CuO_2 layer, as explained above. In Fig.1 we illustrate typical results for the angle-integrated spectral function and a doping $x \sim 20\%$. For comparison in Fig.1(c) the density of states of the non-interacting CuO_2 plane (i.e. with $U_d = 0$) is shown. The bonding-band (mainly copper) is located around $\sim -9\text{eV}$, the antibonding band (mainly oxygen) extends from $\sim -3\text{eV}$ to $\sim 1\text{eV}$ and the non-bonding oxygen orbitals is in between. For every band the typical 2D logarithmic van-Hove singularity is clearly visible, for the non-bonding band, for instance, it is located at $\mu - 5.3\text{eV} = -3.1\text{eV}$. In Fig.1(b) the location and the weight of the peaks of the CuO_4 -cluster Green's function are shown.

The results for the full Green's function of the extended CuO_2 layer are presented in Fig.1(a). Coherently propagating many-body bands have formed out of the cluster energy-levels⁽¹⁸⁾. The chemical potential lies in the band of Zhang-Rice singlets⁽²¹⁾, separated by a charge-transfer gap of about $\sim 1\text{eV}$ from the empty band of copper- d^{10} states. The non-bonding oxygen orbitals and the band of Zhang-Rice triplets are located -3eV to -5.5eV below the Fermi-level.

Van Hove singularities A van Hove singularity can be observed in the band of Zhang-Rice singlets (see Fig.1(a)). It is tempting to identify this structure with the very

sharp feature observed in recent angle-resolved photoemission experiments⁽⁵⁾ on untwinned $\text{YBa}_2\text{Cu}_3\text{O}_{6.9}$ about 1eV below the Fermi-surface. Let us note that, contrary to predictions based on a weak-coupling picture, the van Hove singularity in the density of states does not occur at half-filling in our many body calculation. We find that only for relative large dopings, namely of about 38%, the Fermi-surface would be located right at the van Hove singularity. The absence of a van Hove singularity at the Fermi-edge for dopings $x < 0.2$ is consistent with recent NMR experiments⁽²²⁾ and in contrast to a band-structure calculation based on the local density approximation⁽²³⁾. Also we note that these results do not support the recently proposed “van Hove scenario” of cuprate superconductivity⁽²⁴⁾.

Fermi surface In Fig.2(a) we present the results for the Fermi-surface as a function of doping, $x = 0.01, \dots, 0.70$. Also given are values of the quasiparticle spectral weight, $Z_{\mathbf{k}}$, along the (1,1) direction at the Fermi edge. In Fig.2(b) we present, for comparison, the Fermi-surface of the non-interacting CuO_2 layer at various dopings. (0,0) corresponds to the Γ -point.

We find that the Fermi-surface presented in Fig.2(a) is in agreement with angular resolved photoemission experiments^(1,3) for dopings of about $x \sim 20\%$. The location along the (1,0)—(1,1) lines does also agree quantitatively quite well. Along the (0,0)—(1,1) line the calculated Fermi edge is located at somewhat too large momenta. We note that the quasiparticle spectral weight is reduced by a factor 2 to 5, depending on the doping. For small dopings the calculated Fermi surface expands rapidly, due to the reduced spectral weight, $Z_{\mathbf{k}}$ (compare Eq. (3)) and finally closes completely for $x = 0$ at the (1,1) point. Similar results for the volume of the Fermi-sea as a function of doping has been found by Unger and Fulde⁽¹²⁾ with the projection technique. Note that the data presented in Fig.1 are for a paramagnetic state and are therefore not in contradiction with the notion^(13,25) that in the antiferromagnetic state and dopings of a few percent, the Fermi-surface should take the form of “hole-pockets” centered around $(\pi/2, \pi/2)$.

For small interaction strength, in perturbation theory, the Luttinger sum-rule⁽²⁶⁾ is valid. It states, that the volume of the Fermi-sea of the system with interaction should be equal to that of the non-interacting system. Comparing Fig.2(b) and Fig.2(a) we note that the volume of the Fermi sea obtained for $U_d = 8.8\text{eV}$ is larger than the one predicted by the Luttinger sum-rule (i.e. those obtained for $U_d = 0.0$). E.g. for $x = 20\%$ the difference is $\sim 14\%$ of the Brillouin zone. The cluster-expansion for the self-energy used here is best for large interacting strength and it is therefore not so surprising that the results for the Fermi-surface violate the Luttinger sum-rule, which may be expected to hold for small interactions, U_d . Let us note, that for very large dopings the Fermi-surface for both $U_d = 8.8\text{eV}$ and $U_d = 0$ have nearly identical volumes (for $x = 70\%$ they agree within a percent), since the doped charge carriers go predominantly into the oxygen orbitals, which are influenced only indirectly by U_d .

Quantitatively the Luttinger sum-rule can be experimentally verified best by comparison with results from band-structure calculations. The experimentally determined Fermi-surfaces (accuracy⁽⁴⁾: 10-20% of the Brillouin zone) agree quite well with band-structure

calculations⁽⁷⁾. Since, on the other hand, standard band-structure calculations employ only one Slater determinant they necessarily fulfill the Luttinger sum-rule (and yield a $Z_k \equiv 1$).

Chemical potential In Fig.3 we present the shift (relative to half-filling⁽²⁷⁾) in the chemical potential, μ , upon doping, x . For comparison the shift in the chemical potential for the non-interacting CuO_2 plane, μ_0 , is also shown, they differ by about a factor of two. We have included in Fig.3 the data obtained by a recent photoemission study⁽⁶⁾ on $\text{Bi}_2\text{Sr}_2\text{Ca}_{1-x}\text{Y}_x\text{Cu}_2\text{O}_{8+\delta}$. In order to compare with the experimental results we adjusted our calculated position of the chemical potential for $x = 0$ to the experimental value for $x = 0.01$. Note that, eventhough the experimental data for $x = 0.01 \dots 0.03$ correspond to an insulator state, while we have a metallic state, excluding these points and adjusting at values, e.g., $x = 0.08$, wouldn't change the curves significantly.

We also show in Fig.3 the doping dependence of the renormalized Fermi-velocity, $v_F/v_{F0} \sim m/m^*$. We find, for about $x = 20\%$ doping, an effective mass m^*/m of about two, in agreement with results from angle-resolved photoemission experiments⁽¹⁾ and quantum Monte-Carlo studies⁽¹⁷⁾. The effective mass is found to diverge for dopings $x \rightarrow 0$ (see Fig.3), as a signal of the metal-insulator transition at half filling.

Conclusions It is still controversial how the Fermi surface of 2D correlated, translational invariant electrons on a lattice changes when going from the metal to the Mott-Hubbard insulator regime as a function of electron density. While it seems clear⁽¹³⁾ that the first holes doped into the half-filled, antiferromagnetic ordered insulator, occupy states with momenta around $(\pm\pi/2, \pm\pi/2)$, the situation for finite doping concentrations in the paramagnetic state is not so clear. Here we proposed a scenario, where the volume of the Fermi-sea expands continuously with particle density towards the (π, π) point at half-filling.

For moderate dopings, $x \sim 0.2\%$, this scenario is compatible with exact diagonalization results of one-band models⁽²⁵⁾ and angle-resolved photoemission⁽³⁾ experiments on the cuprate superconductors. For small dopings, $x = 5 - 10\%$ we predict deviations from the Luttinger sum-rule, which might be measurable. The calculated shift in chemical potential associated with the Fermi-surface topology is in reasonable agreement with recent photoemission experiments⁽⁶⁾.

* * *

We would like to thank A. Kampf, R. Manzke, G.A. Sawatzky, W. Weber and W. Wenzel for discussions. This work was supported by the Deutsche Forschungsgemeinschaft and by the Minister für Wissenschaft und Forschung des Landes Nordrhein-Westfalen.

REFERENCES

- [1] C.G. Olson *et al.*, Phys. Rev. B **42**, 381 (1990).
- [2] G. Mante *et al.*, Zeit. Phys. **80**, 181 (1990).
- [3] J.C. Campuzano *et al.*, Phys. Rev. Lett. **64**, 2308 (1990).
- [4] Liu *et al.*, Phys. Rev. B **46**, 11 056 (1992).
- [5] J.G. Tobin *et al.*, Phys. Rev. B **45**, 5563 (1993).
- [6] M.A. van Veenendaal, R. Schlatmann, G.A. Sawatzky, W.A. Groen, Phys. Rev. B **47**, 446 (1993).
- [7] M.S. Hybertsen, L.F. Mattheiss, Phys. Rev. Lett. **60**, 1161 (1988); W.E. Pickett, Rev. Mod. Phys. **61**, 433 (1989); W.E. Pickett, R.E. Cohen, H. Krakauer, Phys. Rev. B **42**, 8764 (1990).
- [8] See Q. Si, J.P. Lu, K. Levin Phys. Rev. B **45**, 4930 (1992) and references therein.
- [9] V.J. Emery, Phys. Rev. Lett. **58**, 2794 (1987).
- [10] H. Matsumoto, M. Sasaki, S. Ishihara, M. Tachiki, Phys. Rev. B **46**, 3009 (1992); M. Sasaki, H. Matsumoto, M. Tachiki, *ibid.*, 3022 (1992).
- [11] J. Schmalian, G. Baumgärtel, K.-H. Bennemann, Phys. Rev. Lett. **68**, 1406 (1992).
- [12] P. Unger, P. Fulde, *Spectral Function of Holes in the Emery Model*, to be published.
- [13] C.L. Kane, P.A. Lee, N. Read, Phys. Rev. B **39**, 6880 (1989); M.D. Johnson, C. Gros, K.J. von Szczepanski, Phys. Rev. B **43**, 11 207 (1991).
- [14] D. Poilblanc, T. Ziman, H.J. Schulz, E. Dagotto, Phys. Rev. B **47**, 14 267 (1993).
- [15] H. Eskes, L.H. Tjeng, G.A. Sawatzky, Phys. Rev. B **41**, 288 (1990); H. Eskes, G.A. Sawatzky, Phys. Rev. B **44**, 9556 (1991).
- [16] Y. Ohta, K. Tsutsui, W. Koshibae, T. Shimozato, S. Maekawa, Phys. Rev. B **46**, 14 022 (1992).
- [17] G. Dopf, J. Wagner, P. Dietrich, A. Muramatsu, W. Hanke, Phys. Rev. Lett. **68**, 2082 (1992).
- [18] C. Gros, R. Valentí, Phys. Rev. B **48**, 418 (1993).
- [19] J. Hubbard, Proc. R. Soc. A **276**, 238 (1963); **277**, 238 (1964); **281**, 401 (1964).
- [20] In hole-notation the orbital energy difference would be $U_d - 5.3\text{eV} = 3.5\text{eV}$.

- [21] F.C. Zhang, T.M. Rice, Phys. Rev. B **37**, 3759 (1988).
- [22] Y.-Q. Song, M.A. Kennard, K.R. Poppelmeier, W.P. Halperin, Phys. Rev. Lett. **70**, 3131 (1993).
- [23] J.-H. Xu, T.J. Waston-Yang, J. Yu, A.J. Freeman, Phys. Lett. A **120**, 489 (1987) predict a van Hove singularity at the Fermi surface for a doping $x = 0.17$ for $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$.
- [24] D.M. News, C.C. Tsuei, P.C. Pattnaik, C.L. Kane, Comm. Cond. Mat. Phys. **15**, 273 (1992); D.M. News, H.R. Krishnamurthy, P.C. Pattnaik, C.C. Tsuei, C.L. Kane, Phys. Rev. Lett. **69**, 1264 (1992).
- [25] W. Stephan, P. Horsch Phys. Rev. Lett. **66**, 2258 (1991).
- [26] J. Luttinger, J. Ward, Phys. Rev. **118**, 1417 (1960); J. Luttinger, Phys. Rev. **119**, 1153 (1961); G. Benefatto, G. Gallavotti, J. Stat. Phys. **59**, 541 (1990).
- [27] The limit $\lim_{T \rightarrow 0} \lim_{x \rightarrow 0} \mu(x, T)$ for the doping and temperature dependent chemical potential is the middle of the charge transfer-gap. Within our zero-temperature formalism the position of the chemical potential at half-filling, $\lim_{x \rightarrow 0} \lim_{T \rightarrow 0} \mu(x, T)$, coincides with the top of the valence band.

Figure captions

Fig.1 (a) A typical result for the angle-integrated spectral weight, for a doping $x=0.2$. The Fermi surface lies in the band of coherently propagating Zhang-Rice singlets. (b) The position and the weight of the poles of the CuO_4 cluster Green's function used for calculating the self-energy. (c) For comparison the density of states of the non-interacting ($U_d = 0$) CuO_2 plane with the three van Hove singularities for the bonding, non-bonding and the anti-bonding band.

Fig.2 (a) Results for the position of the Fermi surface in the first Brillouin zone, as a function of doping, x , and the quasiparticle renormalization factor, Z , at the Fermi edge. Note the strong reduction of Z near the (π, π) point. (b) For comparison the Fermi surface of the non-interacting ($U_d = 0$) CuO_2 plane, which fulfills the Luttinger sum-rule, is shown.

Fig.3 *Filled circles*: Results for the shift in the chemical potential, μ , upon doping, x , relative to half-filling. *Crosses*: Experimental⁽⁶⁾ results for the shift in chemical potential for $\text{Bi}_2\text{Sr}_2\text{Ca}_{1-x}\text{Y}_x\text{Cu}_2\text{O}_{8+\delta}$. *Open circles*: For comparison the shift in μ_0 of the non-interacting, $U_d = 0$, CuO_2 plane. *Filled squares*: The renormalized Fermi velocity, $v_F/v_{F0} \sim m/m^*$. Note the reduction of about two for $x = 0.2$.